

Orbital liquid in three dimensional Mott insulator: $LaTiO_3$

G.Khaliullin^{1,2}, and S.Maekawa²

¹ *Max-Planck-Institut für Festkörperforschung, Heisenbergstr.1, D-70569 Stuttgart, Germany*

² *Institute for Materials Research, Tohoku University, Sendai 980-8577, Japan*

We present a theory of spin and orbital states in Mott insulator $LaTiO_3$. The spin-orbital superexchange interaction between $d^1(t_{2g})$ ions in cubic crystal suffers from a pathological degeneracy of orbital states at classical level. Quantum effects remove this degeneracy and result in the formation of the coherent ground state, in which the orbital moment of t_{2g} level is fully quenched. We find a finite gap for orbital excitations. Such a disordered state of local degrees of freedom on unfrustrated, simple cubic lattice is highly unusual. Orbital liquid state naturally explains observed anomalies of $LaTiO_3$.

PACS numbers: 75.10.-b, 71.27.+a, 75.30.Et, 75.30.Ds

Experiments on transition metal oxides continue to challenge the theory - Keimer et al. [1] have recently reported on dynamical quenching of t_{2g} orbital angular moments in Mott insulator $LaTiO_3$ which is not associated by any detectable orbital/Jahn-Teller ordering. More specifically, magnon spectra are found to fit a spin one-half, nearly isotropic Heisenberg model on cubic lattice. At the same time, anomalous x-ray scattering study has not revealed any evidence of static orbital order. These observations indicate that t_{2g} orbital degrees of freedom of Ti^{3+} ions are mysteriously missing in a low energy states of $LaTiO_3$. Keimer et al. suggested a picture of strongly fluctuating orbitals to reconcile their results.

Dynamical quenching of local orbital moments in a periodic, simple cubic lattice formed by Ti^{3+} ions in $LaTiO_3$ is fascinating. This poses a serious problem for the canonical Goodenough-Kanamori picture of successive orbital and magnetic orderings - the guiding idea in the modern theory of orbitally degenerate transition metal oxides (see for review [2,3]). Loosely speaking, a lifting of the orbital degeneracy without symmetry breaking is formally similar to the spin moment quenching in periodic Kondo lattices. The novelty of $LaTiO_3$ is however that it is an insulator. At the same time, the ideas of quantum disorder due to the weak connectivity or geometrical frustration (like in pyrochlore systems, see [4] for instance) do not apply here, either.

As discussed below, there are several physical reasons for the orbital disorder in $LaTiO_3$, and the key factor is a special symmetry of t_{2g} -superexchange interaction on cubic lattice. This results in a quantum resonance between orbital levels, which removes degeneracy present at classical level by the formation of resonating bonds in orbital sector - a realization of Anderson's resonating valence bonds (RVB) idea [5] in a three dimensional (3D) insulator with help of orbital degrees of freedom. As for spin moments, they show a long-range order as expected on general grounds. However, a staggered moment is considerably reduced, since short-wavelength magnons are actively involved in the local resonance of exchange bonds - in fact, it is the composite spin-orbital excitation which

plays a crucial role in the theory presented.

We begin with discussion of the t_{2g} -superexchange interaction. In terms of fermions $a_{i,\sigma}$, $b_{i,\sigma}$, $c_{i,\sigma}$ corresponding to t_{2g} levels of yz , zx , xy symmetry, respectively, the hopping term in the Hubbard model reads as $-t \left(a_{i,\sigma}^\dagger a_{j,\sigma} + b_{i,\sigma}^\dagger b_{j,\sigma} + \text{H.c.} \right)$ for the bonds along c -direction in a cubic crystal. Correlation energies in doubly occupied virtual states of Ti^{3+} ion are specified as follows: U for electrons on the same orbital; $U' + J_H$, if they occupy different orbitals and form a spin-singlet; and $U' - J_H$ for a spin-triplet charge excitation. The relation $U' = U - 2J_H$ holds in the atomic limit. As $J_H \ll U$ usually, the dominant part of the superexchange interaction in a cubic lattice is:

$$H_{SE} = -\frac{4t^2}{U} + \sum_{\langle ij \rangle} \left(\mathbf{S}_i \mathbf{S}_j + \frac{1}{4} \right) \hat{J}_{ij}^{(\gamma)}. \quad (1)$$

On c -axis bonds the orbital structure of the exchange "constant" is given by (hereafter, the energy scale $4t^2/U$ is used):

$$\hat{J}_{ij}^{(c)} = n_{i,a} n_{j,a} + n_{i,b} n_{j,b} + a_i^\dagger b_i b_j^\dagger a_j + b_i^\dagger a_i a_j^\dagger b_j. \quad (2)$$

Similar expressions can be obtained for $\hat{J}_{ij}^{(a)}$ and $\hat{J}_{ij}^{(b)}$ by replacing orbital fermions a, b in this equation by b, c and c, a pairs, respectively. We notice, that in Eq.(2) and below fermions have only orbital quantum number (a, b, c orbitons), since a spin component of the original fermions is represented in Eq.(1) by spin one-half operators, as usual. We shall focus now on the Hamiltonian (1), and discuss later the effects of finite Hund coupling corrections (of the order of J_H/U).

As evident from Eq.(1) (consider $\langle \mathbf{S}_i \mathbf{S}_j \rangle = -1/4$), the classical Néel state is infinitely degenerate in orbital sector, reflecting that cubic symmetry is respected by the Néel state. This emphasizes a crucial importance of quantum effects, as has been pointed out first in the context of e_g -systems [6]. The bond directional geometry of e_g orbitals offers a solution [7] in that case: e_g orbital frustration is resolved by order from disorder mechanism selecting a particular (directional) orbital configuration, which maximizes the energy gain from quantum




FIG. 1. (i) Spin-orbital resonance between two (left and right) states of the c -bond pair. (ii) Self-consistent Born approximation for the fermionic orbital self-energy. (iii) Diagram contributing to the bond order parameter χ . Solid (wavy) lines represent orbitons (magnons).

spin fluctuations. t_{2g} orbitals are however not bond oriented (they are all planar), and not much spin fluctuation energy can be gained by any pattern of static orbital orderings. The solution of the t_{2g} -problem proposed here is different: that is an idea of $SU(4)$ spin-orbital excitations recently discussed in a context of one dimensional models [8–10].

Reflecting the geometry of t_{2g} orbitals, every bond in Eq.(1) is represented by two equivalent orbitals (such a symmetry is absent for Ising-like e_g orbitals). This is also evident from the representation of Eq.(2) in terms of orbital pseudospin τ_{ab}^i and density $n_{ab}^i = n_{ia} + n_{ib}$ operators acting on the (a, b) -doublet subspace: $\hat{J}_{ij}^{(c)} = 2 \left(\tau_{ab}^i \tau_{ab}^j + \frac{1}{4} n_{ab}^i n_{ab}^j \right)$. Suppose that $n_{ab}^i = 1$; one then clearly observes an orbital $SU(2)$ symmetry of exchange “integrals”. As we know from spin $SU(2) \times$ orbital $SU(2)$ models [8–10], the exchange energy is gained in that case due to the resonance between degenerate local configurations (spin-singlet $\times$ orbital triplet, and spin-triplet $\times$ orbital singlet), and elementary excitations are a mixed spin-orbital $SU(4)$ modes. Fig.1(i) illustrates the idea of correlated spin-orbital fluctuation. In the present problem, spins and orbitals are different: spin sector is effectively half-filled ($n_{i,\uparrow} + n_{i,\downarrow} = 1$) and must show a long-range order in 3D; on the other hand, the orbital occupation number n_{ab}^i is not conserved but rather fluctuates around the average value $2/3$. Because of the presence of third orbital, one has a less severe constraint $n_{i,a} + n_{i,b} + n_{i,c} = 1$ in orbital sector. Even though a true $SU(4)$ excitation cannot develop here because of spin-orbital asymmetry, the analogy with 1D models suggests the way how to optimize the exchange energy: That is by the formation of a virtual $SU(4)$ resonance which can be viewed in the present context as a local excitation composed of magnon and orbital fluctuation. This resonance removes the orbital frustration dynamically, and the dis-

ordered state of orbitals is precisely what is required to amplify this mechanism in all three directions of cubic lattice.

Technically, the main ingredient of the theory is the three-particle bond variable, $\hat{\chi}_{\alpha}^{(ij)} = \alpha_i^{\dagger} \alpha_j (s_i^{\dagger} + s_j)$. Here, α denotes an appropriate fermionic orbiton, say a or b orbitons for c -bond pair (ij) , while $s^{\dagger}$ being a magnon excitation about the Néel state. This is an analogy to $SU(4)$ flavor bond variable [8] in a spin ordered state. The bare magnon dispersion is simply given by the Heisenberg interaction with an average exchange constant $J = \langle \hat{J}_{ij}^{(\gamma)} \rangle$, which will be calculated below. On the contrary, the orbital dynamics is exclusively due to the three-particle resonance, since spin-only prefactor $\langle \mathbf{S}_i \mathbf{S}_j + 1/4 \rangle$ is almost zero (~ -0.05) in 3D. We may present Eq.(1) as $H_{SE} = H_{sp} + H_{int} + \text{const.}$, where H_{sp} describes bare magnons, and $H_{int} = -\frac{1}{2} \sum_{\langle ij \rangle} \sum_{\alpha\beta} \hat{\chi}_{\alpha}^{(ij)} \hat{\chi}_{\beta}^{(ji)}$. The strategy is then to decouple H_{int} in terms of bond variables. The solution with uniform bond amplitude χ gives $H_{int} \Rightarrow H_a + H_b + H_c + \text{const.}$, where

$$H_a = -\chi \cdot \sum_{\langle ij \rangle_{bc}} \left\{ a_i^{\dagger} a_j (s_i^{\dagger} + s_j) + \text{H.c.} \right\}. \quad (3)$$

Here the summation is restricted to (b, c) -planes. Similar expressions hold for b and c orbitons. Since the filling factor is only $1/3$ per orbital flavor, the orbital sector is far from nesting conditions, thus a mean-field uniform solution in the spirit of large N -theories [11] is believed to be a reasonable starting point to describe a disordered state of orbitals. Eq.(3) describes an effective hopping of orbitons accompanied by simultaneous magnon excitations. Thus, we have mapped the superexchange in an undoped system to an effective $\chi - J$ model, which in a momentum space reads as

$$H_{\chi-J} = \sum_{\mathbf{k}\mathbf{p}} \sum_{\alpha} M_{\mathbf{k}\mathbf{p}}^{(\alpha)} (\alpha_{\mathbf{k}+\mathbf{p}}^{\dagger} \alpha_{\mathbf{k}} s_{\mathbf{p}} + \text{H.c.}) + \sum_{\mathbf{p}} \omega_{\mathbf{p}} s_{\mathbf{p}}^{\dagger} s_{\mathbf{p}}. \quad (4)$$

Here $M_{\mathbf{k}\mathbf{p}}^{\alpha} = -4\chi(\gamma_{\mathbf{k}+\mathbf{p}}^{(\alpha)} u_{\mathbf{p}} + \gamma_{\mathbf{k}}^{(\alpha)} v_{\mathbf{p}})$, while form-factors are $\gamma_{\mathbf{k}}^{(a)} = (c_y + c_z)/2$, $\gamma_{\mathbf{k}}^{(b)} = (c_z + c_x)/2$, $\gamma_{\mathbf{k}}^{(c)} = (c_x + c_y)/2$, where $c_{\lambda} = \cos k_{\lambda}$. Bare magnon dispersion $\omega_{\mathbf{p}} = 3J(1 - \gamma_{\mathbf{p}}^2)^{1/2}$ with $\gamma_{\mathbf{p}} = (c_x + c_y + c_z)/3$, and $u_{\mathbf{p}}, v_{\mathbf{p}}$ are conventional Bogoliubov transformation coefficients for magnons in the cubic lattice. It is noticed that the coupling constant in Eq.(4) vanishes at $p = 0$ limit (spin conservation is respected), and it is the short-wavelength magnons which are important. The model is similar to the $t - J$ model for the doped Néel state of cuprates (see [12,13] for comparison), yet the parameters χ and J have to be self-consistently determined. It is noticed, that there are three branches of 2D fermions, and the dimensionality of magnon sector is three in the present case.

We calculate first the fermionic spectrum within a self-consistent Born approximation. Different from holes in the $t - J$ model, an orbiton creates a magnon on the site it arrives at, and eliminates a magnon while leaving the site (see Eq.(3)). Therefore orbiton motion contains a coherent component even in the Ising limit for magnons. The latter approximation is used to simplify a momentum integration in the self-energy, Fig.1(ii). One then obtains $Re\Sigma_\omega^{(a)}(\mathbf{k}) = f_\omega - \kappa_\omega(\gamma_\mathbf{k}^{(a)})^2$. Here $\kappa_\omega = (4\chi)^2 \int_0^\infty d\xi \rho(\xi)/(\xi + 3J - \omega)$, where $\rho(\xi) = \langle \rho_\mathbf{k}(\xi) \rangle_\mathbf{k}$ is fermionic density of states (DOS). f_ω has similar structure: $f_\omega = (4\chi)^2 \int_{-\infty}^0 d\xi \tilde{\rho}(\xi)/(\omega + 3J - \xi)$, where $\tilde{\rho}(\xi) = \langle \rho_\mathbf{k}(\xi)(\gamma_\mathbf{k}^{(a)})^2 \rangle_\mathbf{k}$. One observes a two-sublattice structure of the orbiton dispersion (imposed by spin order) as expected from analogy with the $t - J$ model [13]. We estimate the fermionic (unrenormalized) mass as $m \simeq 1/\kappa_0$, and density of states as $\rho(0) \simeq 1/\pi\kappa_0$. Each orbiton forms its own 2D fermi-surface (FS). Say, for a orbitons the FS consists of two almost circles around $(0,0)$ and (π,π) points in a (k_y, k_z) plane. Assuming constant DOS within the interval of width $W_{orb} = 1/\rho(0)$, we find $\kappa_0 \simeq 4\chi/\sqrt{\pi}$, and $W_{orb} \simeq 4\sqrt{\pi}\chi$. W_{orb} is the energy scale of orbital fluctuations [14]. Further, we estimate the bond amplitude $\chi = \langle a_i^\dagger a_j (s_i^\dagger + s_j) \rangle$. In the Ising limit for intermediate magnons, Fig.1(iii) gives $\chi \simeq (4/3\sqrt{\pi}) \langle (\gamma_\mathbf{k}^{(a)})^2 \rangle_{FS}$. For $\langle n^{(a)} \rangle = 1/3$ we estimate $\langle (\gamma_\mathbf{k}^{(a)})^2 \rangle_{FS} \simeq 0.3$. Spin stiffness in the present model is controlled by J . The mean-field value $2/9$ for J follows from Eq.(2). This number is reduced to $J \simeq 0.16$ due to the corrections shown in Fig.2(i). The physics behind this reduction is that $SU(4)$ resonance induces some ferromagnetic component in spin interactions.

Thus, we have fixed basic energy scales (in units of $4t^2/U$): $W_{orb} \simeq 1.6$ for orbital sector, and $3J \simeq 0.48$ for magnon bandwidth. The spin stiffness is relatively small because Pauli principle is relevant only if electrons occupy the same orbital, and that probability is reduced at large orbital degeneracy. Main energy gain stems from magnon-orbital resonance: $E_{int} = -6\chi^2 \simeq -0.31$. (Spin-only contribution is negligible: $E_{SW} \simeq -0.02$). This number should be compared with energy gain that one has with statically ordered planar orbitals: $\simeq -0.17$.

We discuss now physical consequences of the theory in context of experimental data on $LaTiO_3$ [1]. Observed magnon dispersions of cubic symmetry are very natural in above picture of orbitals fluctuating faster than magnons. The energy scale $4t^2/U \simeq 100$ meV, required to fit $J_{exp} = 15.5 \pm 1$ meV, is also reasonable in view of values $U \sim 4$ eV [15], and $t \sim 0.3$ eV [16]. This scale means that orbital fluctuation energy in $LaTiO_3$ is $W_{orb} \sim 160$ meV.

Keimer et al. observed very small anisotropy gap, indicating that effects of conventional relativistic spin-orbit coupling are strongly suppressed. This is in fact an un-




FIG. 2. (i) Interaction corrections to the spin exchange constant J . (ii) Magnon self-energies.

avoidable consequence of our theory, since the ground state has no orbital degeneracy. We may calculate spectral density of orbital angular momentum fluctuations. It is given by the fermionic orbiton excitations (“orbiton Stoner continuum”). Spectral density of the local susceptibility of angular moments $l_i^z = i(a_i^\dagger b - b^\dagger a)_i$ vanishes in the static, $\omega = 0$ limit: $S(\omega) = 2\rho^2(0)\omega$. This implies complete quenching of orbital moments at low energies. Making parallel with Jahn-Teller (JT) impurity physics [17], we may say that “effective Ham reduction factor”, $\zeta(\bar{\omega}) \sim \left\{ \int_0^{\bar{\omega}} S(\omega) d\omega \right\}^{1/2} \sim \bar{\omega}/W_{orb}$ is frequency dependent, and the angular moments disappear from low-energy physics linearly in energy. Also, the ratio $\Lambda_{so}/W_{orb} \sim 10^{-1}$ ($\Lambda_{so} \sim 20$ meV [1]) means that relativistic spin-orbit coupling induced corrections are small; we estimate the g -value shift $\Delta g/g \sim 0.06$, consistent with [1].

Despite three dimensionality of magnon spectra, the staggered moment in $LaTiO_3$ is small, i.e. $0.45\mu_B$ [1] - an obvious problem for a spin-wave picture. The present theory resolves this difficulty: The intensity of spin Bragg peak is partially taken away by a quantum magnon-orbital resonance and redistributed over finite frequency region. In other words, fluctuating orbitals generate additional quantum spin fluctuations in the ground state. We have calculated a spin moment reduction in a similar way as it was previously done in the $t - J$ model [18]. Accounting for lowest order interaction corrections (Fig.2(ii)) to the magnons, one finds:

$$\delta S_{int}^z = \frac{3}{2}(4\chi)^2 \sum_{\mathbf{k}\mathbf{k}'} n_{\mathbf{k}}(1 - n_{\mathbf{k}'}) \left\{ \frac{A_{\mathbf{k}\mathbf{k}'}}{(\omega_{\mathbf{p}} + \xi_{\mathbf{k}'} - \xi_{\mathbf{k}})^2} - \frac{B_{\mathbf{k}\mathbf{k}'}}{\omega_{\mathbf{p}}(\omega_{\mathbf{p}} + \xi_{\mathbf{k}'} - \xi_{\mathbf{k}})} \right\}, \quad (5)$$

where $\xi_{\mathbf{k}}$ is the a orbiton dispersion, $\mathbf{k}' = \mathbf{k} + \mathbf{p}$, $A_{\mathbf{k}\mathbf{k}'} = \lambda_+^2/(1 + \gamma_{\mathbf{p}}) + \lambda_- \lambda_-/(1 - \gamma_{\mathbf{p}}^2)^{1/2}$, $B_{\mathbf{k}\mathbf{k}'} = \lambda_+^2 \gamma_{\mathbf{p}}/(1 + \gamma_{\mathbf{p}})$, and $\lambda_{\pm} = \gamma_{\mathbf{k}}^{(a)} \pm \gamma_{\mathbf{k}'}^{(a)}$. By averaging first the matrix elements A, B in Eq.(5) over the Brillouin zone (they are rather regular functions), we evaluate $\delta S_{int}^z \simeq 0.15$. (Another estimation by using the Ising limit for magnons gives 0.13). Adding also a conventional 3D spin-wave

correction $\delta S_{SW}^z \simeq 0.075$, one then obtains a staggered moment $0.55\mu_B$ in fair agreement with experiment.

The orbital liquid picture offers a simple explanation of puzzling Fano-type phonon anomalies observed by Raman scattering data in insulating titanates (Reedyk et al. [19]). These anomalies are most pronounced in $LaTiO_3$, and we identify their origin as due to the coupling of phonons to orbital excitations, noticing that the phonon position ($\sim 300 \text{ cm}^{-1}$) is right in the orbital Stoner continuum.

According to Goodenough-Kanamori rules, a local ferrottype orbital correlations are expected in a spin Néel state. A residual interactions (via magnons) between orbitons do indeed produce a triplet pairing of orbitons within each fermionic branch. Treating the problem on BCS level, we find finite mean-field order parameter $\langle \alpha_i^\dagger \alpha_j^\dagger \rangle$ of p-wave symmetry (specifically, of $p_x + ip_y$ -symmetry for c orbitons, for instance), which opens an orbital gap. It is difficult to calculate reliably the value of this gap (of the order of a few meV); we predict however that a linear γ -term in specific heat should be released when a correlation gap in orbital spectrum is thermally washed out. We estimate $\gamma \sim 40 \text{ mJ/moleK}^2$. Specific heat measurements in $LaTiO_3$ up to $T \sim 100 \text{ K}$ would be a crucial test for the theory.

From physical point of view, the proximity of $LaTiO_3$ to Mott transition (charge gap is only 0.2 eV [16]) is actually a key factor, which drives this compound to superexchange dominated orbital liquid state. This state is stable, however, only if the effective orbiton bandwidth, $W_{orb} \sim 4t^2/U$, is large enough to suppress JT order. When one goes to another end compound, $YTiO_3$ (charge gap $\sim 1 \text{ eV}$) the bandwidth narrows [16], thus reducing superexchange energy scale, and we expect static JT ordering of spatially more localized orbitals once the ratio W_{orb}/E_{JT} becomes less than critical. With antiferrottype orbital ordering as observed in $YTiO_3$ [20] a classical expectation value of Eq.(2) vanishes, and Hund's coupling J_H term induces a ferromagnetic state. The competition between the frustrated superexchange on one hand, and JT plus Hund's couplings on the other hand is expected to result in the phase diagram shown in Fig.3.

To conclude, we have argued that due to large degeneracy and special geometry of orbitals t_{2g} -superexchange system most likely has an orbitally disordered ground state. The observed anomalies of $LaTiO_3$ find their natural explanations in a proposed orbital liquid picture. In metallic manganites, e_g orbital disorder is enforced by the presence of mobile carriers [21]. Apparently, the case of insulating $LaTiO_3$ is a nice example of that a frustrated t_{2g} orbitals can do the same job alone, without doping and well before the Mott transition to a metallic state is reached.

We would like to thank B. Keimer for stimulating discussions. Discussions with Y. Tokura, S. Ishihara, P. Prelovšek, and K. Tsutsui are also acknowledged. This




FIG. 3. Proposed phase diagram for $d^1(t_{2g})$ Mott insulator on cubic lattice. At small Hund's and Jahn-Teller couplings the quantum Néel state with dynamically quenched orbital moments is stabilized. Quantum phase transition line separates this state from the ferromagnetic phase with static orbital/JT order.

work was supported by Priority Areas Grants from the Ministry of Education, Science, Culture and Sport of Japan, CREST and NEDO.

-
- [1] B. Keimer *et al.*, (preprint cond-mat/0002014).
 - [2] Y. Tokura, and N. Nagaosa, *Science* **288**, 462 (2000).
 - [3] K. I. Kugel, and D. I. Khomskii, *Sov.Phys.Usp.* **25**, 231 (1982).
 - [4] B. Canals, and C. Lacroix, *Phys. Rev. Lett.* **80**, 2933 (1998).
 - [5] P. W. Anderson, *Science* **235**, 1196 (1987).
 - [6] L. F. Feiner, A. M. Oleś, and J. Zaanen, *Phys. Rev. Lett.* **78**, 2799 (1997).
 - [7] G. Khaliullin, and V. Oudovenko, *Phys. Rev. B* **56**, R14243 (1997); G. Khaliullin, and R. Kilian, *J. Phys.: Condens. Matter*, **11**, 9757 (1999).
 - [8] Y. Q. Li, M. Ma, D. N. Shi, and F. C. Zhang, *Phys. Rev. Lett.* **81**, 3527 (1998).
 - [9] B. Frischmuth, F. Mila, and M. Troyer, *Phys. Rev. Lett.* **82**, 835 (1999).
 - [10] F. Mila, B. Frischmuth, A. Deppeler, and M. Troyer, *Phys. Rev. Lett.* **82**, 3697 (1999).
 - [11] I. Affleck, and J. B. Marston, *Phys. Rev. B* **37**, 3774 (1988).
 - [12] C. L. Kane, P. A. Lee, and N. Read, *Phys. Rev. B* **39**, 6880 (1989).
 - [13] G. Martinez, and P. Horsch, *Phys. Rev. B* **44**, 317 (1991).
 - [14] More specifically, orbital dynamics contains both coherent and incoherent components. With χ, J given in the text, we have estimated the quasiparticle weight ~ 0.7 , which is close to the ratio J/χ .
 - [15] T. Mizokawa, and A. Fujimori, *Phys. Rev. B* **54**, 5368 (1996); T. Mizokawa, D. I. Khomskii, and G. A. Sawatzky, *Phys. Rev. B* **60**, 7309 (1999).
 - [16] T. Katsufuji, Y. Taguchi, and Y. Tokura, *Phys. Rev. B* **56**, 10145 (1993).
 - [17] F. S. Ham, *Phys. Rev.* **138**, A1727 (1965); **166**, 307 (1968).

- [18] G. Khaliullin, and P. Horsch, Phys. Rev. B **47**, 463 (1993).
- [19] M. Reedyk *et al.*, Phys. Rev. B **55**, 1442 (1997).
- [20] H. Nakao et al., to be published
- [21] S. Ishihara, M. Yamanaka, and N. Nagaosa, Phys. Rev. B **56**, 686 (1997); R. Kilian, and G. Khaliullin, Phys. Rev. B **58**, R11841 (1998).